



On equitable coloring of corona of wheels

J. Vernold Vivin^a, K. Kaliraj^b

^aDepartment of Mathematics, University College of Engineering Nagercoil, (Anna University Constituent College), Konam, Nagercoil- 629 004, Tamil Nadu, India.

^bDepartment of Mathematics, Ramanujan Institute for Advanced Study in Mathematics, University of Madras, Chepauk, Chennai-600 005, Tamil Nadu, India.

vernoldvivin@yahoo.in, sk.kaliraj@gmail.com

Abstract

The notion of equitable colorability was introduced by Meyer in 1973 [9]. In this paper we obtain interesting results regarding the equitable chromatic number $\chi_{=}$ for the corona graph of a simple graph with a wheel graph $G \circ W_n$. Some extensions into l -corona products are also determined.

Keywords: equitable coloring, corona graph, wheel graph

Mathematics Subject Classification : 05C15

DOI:10.5614/ejgta.2016.4.2.8

1. Introduction

If the set of vertices of a graph G can be partitioned into k classes V_1, V_2, \dots, V_k such that each V_i is an independent set and the condition $||V_i| - |V_j|| \leq 1$ holds for every pair (i, j) , then G is said to be *equitably k -colorable*. The smallest integer k for which G is equitably k -colorable is known as the *equitable chromatic number* [9] of G and denoted by $\chi_{=}(G)$. This subject is widely discussed in literature [1, 4, 6, 7, 9]. In general, the problem of optimal equitable coloring, in the sense of number color used, is NP-hard.

The *corona* of two graphs G_1 and G_2 is the graph $G = G_1 \circ G_2$ formed from one copy of G_1 and $|V(G_1)|$ copies of G_2 where the i th vertex of G_1 is adjacent to every vertex in the i th copy of G_2 . For any integer $l \geq 2$, we define the graph $G_1 \circ^l G_2$ recursively from $G_1 \circ G_2$ as

Received: 9 February 2015, Revised: 18 September 2016, Accepted: 29 September 2016.

$G_1 \circ^l G_2 = (G_1 \circ^{l-1} G_2) \circ G_2$. Graph $G_1 \circ^l G_2$ is also named as l -corona product of G_1 and G_2 . This kind of product was introduced by Harary and Frucht in 1970 [2].

Even more, we know [4] that the problem of the equitable coloring of corona graphs $G \circ H$ is NP-hard when G is 4-regular graph and $H = K_2$. So we have to look for simplified structure of graphs allowing polynomial-time algorithms. This paper gives such solutions for corona graph of a simple graph with a wheel graph. Some extensions for l -corona products are also determined. This way we confirm Equitable Coloring Conjecture posed by Meyer [9] for graphs under consideration.

Conjecture 1 (Equitable Coloring Conjecture (ECC) [9]). *For any connected graph G , other than the complete graph or odd cycle, $\chi_=(G) \leq \Delta(G)$.*

This conjecture has been verified for all graphs with six or fewer vertices. Lih and Wu [7] proved that the Equitable Coloring Conjecture (ECC) is true for all bipartite graphs. Wang and Zhang [10] considered a broader class of graphs, namely r -partite graphs. They proved that Meyer’s conjecture is true for complete graphs from this class. The conjecture (or even the stronger one) was confirmed for outerplanar graphs [11] and planar graphs with maximum degree at least 13 [12].

Graph products are interesting and useful in many situations [5]. Equitable coloring of Cartesian, weak tensor and strong tensor products for some classes of graphs was considered in [3, 8].

For any integer $n \geq 4$, the *wheel graph* W_n is the n -vertex graph obtained by joining a vertex v_1 to each of the $n - 1$ vertices $\{w_1, w_2, \dots, w_{n-1}\}$ of the cycle graph C_{n-1} .

2. Equitable coloring on corona graph of simple graph with wheel graph

We start with giving results for coronas of a single vertex and a wheel.

Theorem 2.1. *Let n be a positive integer, $n \geq 4$. Then*

$$\chi_=(K_1 \circ W_n) = \left\lceil \frac{n-1}{2} \right\rceil + 2.$$

Proof. The color used for coloring the vertex of K_1 and the color used for coloring vertex v_1 cannot be used more times, so we can use any other color at most twice. Hence the value of the equitable chromatic number is equal to $\left\lceil \frac{n-1}{2} \right\rceil + 2$. □

We notice that $\Delta(K_1 \circ W_n) = n \geq \lceil (n-1)/2 \rceil + 2$ for $n \geq 4$. This means that ECC holds for $K_1 \circ W_n, n \geq 4$.

Next, we consider coronas, where the set of vertices of graph G includes more than one element.

Theorem 2.2. *Let G be an equitably 4-colorable graph on, $m \geq 2$, vertices and let m be even, n be odd, and $n \geq 4$, then*

$$\chi_=(G \circ W_n) = 4.$$

Proof. Let $n_i(k)$ be the number of appearance of color k , $1 \leq k \leq 4$, in the i th copy of W_n corresponding to vertex u_i of G in $G \circ W_n$, $i = 1, 2, \dots, m$.

Let $f(u_i) = j$ be the color assigned to vertex u_i ($1 \leq i \leq m$) of G . Since G is 4-colorable j takes the values in the range $1 \leq j \leq 4$.

We color graph G equitably with four colors. We order the vertices of G : u_1, u_2, \dots, u_m in such a way that vertex u_i is colored with color $i \bmod 4$ - we use color 4 instead of color 0 (in some cases recoloring is needed). We extend this coloring into whole graph $G \circ W_n$ due to the following conditions. We consider two cases:

1. $m \bmod 4 \equiv 0$

If $f(u_i) = j$, $u_i \in V(G)$, $1 \leq j \leq 4$, then

- $n_i((j + 1) \bmod 4) = 1$,
- $n_i((j + 2) \bmod 4) = \frac{n - 1}{2}$ and
- $n_i((j + 3) \bmod 4) = \frac{n - 1}{2}$.

In the above coloring, we use each color exactly $(n + 1)m/4$ times. Graph $G \circ W_n$ is colored equitably.

2. $m \bmod 4 \equiv 2$

We color first $m - 2$ copies of W_n as we have colored the corresponding vertices in Case (1). We color last two copies in the following way. For each vertex u_i , $i = m - 1, m$, if $f(u_i) = j$, $1 \leq j \leq 2$, then the extended coloring must fulfill the following conditions.

- $n_i((j + 2)) = 1$,
- $n_i((j + 3) \bmod 4) = \frac{n - 1}{2}$,
- $n_i(j + 1) = \frac{n - 1}{2}$.

We use each of four colors exactly $(n + 1)\lfloor m/4 \rfloor + (n + 1)/2$ times. Graph $G \circ W_n$ is colored equitably.

Hence $\chi_=(G \circ W_n) \leq 4$. By the definition of corona graph, graph $G \circ W_n$ contains K_4 . Hence $\chi_=(G \circ W_n) = 4$. □

Theorem 2.3. *Let G be an equitably 4-colorable graph on 5 vertices, then*

$$\chi_=(G \circ W_5) = 4.$$

Proof. Since W_5 has the cycle C_4 , $\chi(W_4) \geq 3$. By the definition of corona, each vertex u_i of G is adjacent to every vertex of its copy of W_n . Hence $\chi_=(G \circ W_5) \geq 4$.

By assigning the colors 1,2,3 and 4 as given below, it is concluded that the 1 appears 7 times, 2 appears 8 times, 3 appears 8 times and 4 appears 7 times. (i.e) The difference between the number of appearance of each pair of colors does not exceed one. Hence $\chi_=(G \circ W_5) \leq 4$. Hence $\chi_=(G \circ W_5) = 4$. □




Figure 1. An equitable 4-coloring of $K_{1,1,1,2} \circ W_5$ with $n(1) = n(4) = 7$ and $n(2) = n(3) = 8$.

Now, we consider the remaining cases of m and n . It turns out that in these cases five colors are desirable for proper equitable coloring.

Theorem 2.4. *Let G be an equitably 5-colorable graph on m vertices. If $m \bmod 2 \equiv 1, n \geq 7$ or m, n even with $n \geq 4$ then*

$$\chi_=(G \circ W_n) = 5.$$

Proof. Let $n_i(k)$ be the number of appearance of color $k, 1 \leq k \leq 5$, in the i th copy of W_n corresponding to vertex u_i of G in $G \circ W_n, i = 1, 2, \dots, m$.

Let $f(u_i) = j$ be the color assigned to vertex $u_i (1 \leq i \leq m)$ of G . Since G is 5-colorable j takes the values in the range $1 \leq j \leq 5$.

We color graph G equitably with five colors. We order the vertices of $G: u_1, u_2, \dots, u_m$ in such a way that vertex u_i is colored with color $i \bmod 5$ - we use color 5 instead of color 0 (in some cases recoloring is needed). We extend this coloring to the whole graph $G \circ W_n$ due to the following conditions. We consider five cases dependently on the value of m .

1. $m \bmod 5 \equiv 0$

For each vertex $u_i \in V(G)$ if $f(u_i) = j, 1 \leq j \leq 5$, then

- $n_i((j + 1) \bmod 5) = 1,$
- $n_i((j + 2) \bmod 5) = 1,$
- $n_i((j + 3) \bmod 5) = \left\lceil \frac{n - 2}{2} \right\rceil,$

- $n_i((j + 4) \bmod 5) = \left\lfloor \frac{n - 2}{2} \right\rfloor$

We use each of the five colors exactly $(n + 1)m/5$ times. Graph $G \circ W_n$ is colored equitably.

2. $m \bmod 5 \equiv 1$

First, we color $m - 6$ copies of W_n as we color the corresponding vertices in Case (1). We color last six copies in the following way. For each vertex u_i ($m - 5 \leq i \leq m$) we extend the coloring due to the following conditions, dependently on n .

(a) $n \bmod 5 \equiv 0$

- For vertex u_{m-5} ($f(u_{m-5}) = 1$) we have $n_{m-5}(2) = 1, n_{m-5}(3) = n_{m-5}(4) = \frac{2n - 5}{5}, n_{m-5}(5) = \frac{n + 5}{5}$.
- For vertex u_{m-4} ($f(u_{m-4}) = 2$) we have $n_{m-4}(3) = 1, n_{m-4}(1) = n_{m-4}(5) = \frac{2n - 5}{5}, n_{m-4}(4) = \frac{n + 5}{5}$.
- For vertex u_{m-3} ($f(u_{m-3}) = 3$) we have $n_{m-3}(4) = 1, n_{m-3}(2) = \frac{2n}{5}, n_{m-3}(5) = \frac{2n - 10}{5}, n_{m-3}(1) = \frac{n + 5}{5}$.
- For vertex u_{m-2} ($f(u_{m-2}) = 4$) we have $n_{m-2}(5) = 1, n_{m-2}(3) = \frac{2n}{5}, n_{m-2}(2) = \frac{2n - 5}{5}, n_{m-2}(1) = \frac{n}{5}$.
- For vertex u_{m-1} ($f(u_{m-1}) = 5$) we have $n_{m-1}(3) = 1, n_{m-1}(1) = n_{m-1}(2) = \frac{2n - 5}{5}, n_{m-1}(4) = \frac{n + 5}{5}$.
- For vertex u_m ($f(u_m) = 1$) we have $n_m(2) = 1, n_m(3) = n_m(4) = \frac{2n - 5}{5}, n_m(5) = \frac{n + 5}{5}$.

Each of the colors 1, 2, 3 and 5 are used $(6n + 5)/5$ times and color 4 is used $(6n + 5)/5 + 1$ times.

(b) $n \bmod 5 \equiv 1$ or $n \bmod 5 \equiv 4$

- For vertex u_{m-5} ($f(u_{m-5}) = 1$) we have $n_{m-5}(2) = 1, n_{m-5}(3) = n_{m-5}(4) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-5}(5) = \left\lfloor \frac{n - 1}{5} \right\rfloor$.
- For vertex u_{m-4} ($f(u_{m-4}) = 2$) we have $n_{m-4}(3) = 1, n_{m-4}(1) = n_{m-4}(5) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-4}(4) = \left\lfloor \frac{n - 1}{5} \right\rfloor$.
- For vertex u_{m-3} ($f(u_{m-3}) = 3$) we have $n_{m-3}(4) = 1, n_{m-3}(2) = n_{m-3}(5) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-3}(1) = \left\lfloor \frac{n - 1}{5} \right\rfloor$.
- For vertex u_{m-2} ($f(u_{m-2}) = 4$) we have $n_{m-2}(5) = 1, n_{m-2}(2) = n_{m-2}(3) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-2}(1) = \left\lfloor \frac{n - 1}{5} \right\rfloor$.

- For vertex u_{m-1} ($f(u_{m-1}) = 5$) we have $n_{m-1}(3) = 1, n_{m-1}(1) = n_{m-1}(2) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-1}(4) = \left\lfloor \frac{n-1}{5} \right\rfloor$.
- For vertex u_m ($f(u_m) = 1$) we have $n_m(2) = 1, n_m(3) = n_m(4) = \left\lfloor \frac{2n}{5} \right\rfloor, n_m(5) = \left\lfloor \frac{n-1}{5} \right\rfloor$.

Each of the colors 1, 4 and 5 are used $2 + 2\lfloor 2n/5 \rfloor + 2\lfloor (n-1)/5 \rfloor$ times and colors 2 and 3 are used, each one with, $3 + 3\lfloor 2n/5 \rfloor$ times. For $n \bmod 5 \equiv 1$ or $n \bmod 5 \equiv 4$, the difference does not exceed one.

(c) $n \bmod 5 \equiv 2$

- For vertex u_{m-5} ($f(u_{m-5}) = 1$) we have $n_{m-5}(2) = 1, n_{m-5}(3) = \frac{2n+1}{5}, n_{m-5}(4) = \frac{2n-4}{5}, n_{m-5}(5) = \frac{n-2}{5}$.
- For vertex u_{m-4} ($f(u_{m-4}) = 2$) we have $n_{m-4}(3) = 1, n_{m-4}(1) = \frac{2n-4}{5}, n_{m-4}(5) = \frac{2n+1}{5}, n_{m-4}(4) = \frac{n-2}{5}$.
- For vertex u_{m-3} ($f(u_{m-3}) = 3$) we have $n_{m-3}(4) = 1, n_{m-3}(2) = \frac{2n+1}{5}, n_{m-3}(5) = \frac{2n-4}{5}, n_{m-3}(1) = \frac{n-2}{5}$.
- For vertex u_{m-2} ($f(u_{m-2}) = 4$) we have $n_{m-2}(5) = 1, n_{m-2}(2) = \frac{2n+1}{5}, n_{m-2}(3) = \frac{2n-9}{5}, n_{m-2}(1) = \frac{n+3}{5}$.
- For vertex u_{m-1} ($f(u_{m-1}) = 5$) we have $n_{m-1}(3) = 1, n_{m-1}(1) = \frac{2n+1}{5}, n_{m-1}(2) = \frac{2n-9}{5}, n_{m-1}(4) = \frac{n+3}{5}$.
- For vertex u_m ($f(u_m) = 1$) we have $n_m(2) = 1, n_m(3) = \frac{2n+1}{5}, n_m(4) = \frac{2n-4}{5}, n_m(5) = \frac{n-2}{5}$.

Each of the colors 1, 2 and 3 are used $(6n+3)/5 + 1$ times and colors 4 and 5 are used, each one with, $(6n+3)/5$ times.

(d) $n \bmod 5 \equiv 3$

- For vertex u_{m-5} ($f(u_{m-5}) = 1$) we have $n_{m-5}(2) = 1, n_{m-5}(3) = n_{m-5}(4) = \frac{2n-1}{5}, n_{m-5}(5) = \frac{n-3}{5}$.
- For vertex u_{m-4} ($f(u_{m-4}) = 2$) we have $n_{m-4}(3) = 1, n_{m-4}(1) = n_{m-4}(5) = \frac{2n-1}{5}, n_{m-4}(4) = \frac{n-3}{5}$.

- For vertex u_{m-3} ($f(u_{m-3}) = 3$) we have $n_{m-3}(4) = 1, n_{m-3}(2) = n_{m-3}(5) = \frac{2n-1}{5}, n_{m-3}(1) = \frac{n-3}{5}$.
- For vertex u_{m-2} ($f(u_{m-2}) = 4$) we have $n_{m-2}(5) = 1, n_{m-2}(2) = \frac{2n-1}{5}, n_{m-2}(3) = \frac{2n-6}{5}, n_{m-2}(1) = \frac{n+2}{5}$.
- For vertex u_{m-1} ($f(u_{m-1}) = 5$) we have $n_{m-1}(3) = 1, n_{m-1}(1) = \frac{2n-1}{5}, n_{m-1}(2) = \frac{2n-6}{5}, n_{m-1}(4) = \frac{n+2}{5}$.
- For vertex u_m ($f(u_m) = 1$) we have $n_m(2) = 1, n_m(3) = n_m(4) = \frac{2n-1}{5}, n_m(5) = \frac{n-3}{5}$.

Each of the colors 1, 2, 3 and 4 are used $(6n+2)/5 + 1$ times and color 5 is used $(6n+2)/5$ times.

In all the above cases the difference between the cardinalities of the color classes does not exceed one, so our coloring is equitable.

3. $m \bmod 5 \equiv 2$

We color first $m-2$ copies of W_n as we color the corresponding vertices in Case (1). We color last two copies (for u_{m-1} and u_m) in the following way. We consider five cases dependently on n .

(a) $n \bmod 5 \equiv 0$

- If $f(u_i) = 1, n_i(3) = 1, n_i(2) = \frac{2n-5}{5}, n_i(4) = \frac{2n-5}{5}, n_i(5) = \frac{n+5}{5}$.
- If $f(u_i) = 2, n_i(4) = 1, n_i(1) = n_i(3) = \frac{2n}{5}, n_i(5) = \frac{n-5}{5}$.

(b) $n \bmod 5 \equiv 1$ or $n \bmod 5 \equiv 4$

- If $f(u_i) = 1, n_i(3) = 1, n_i(2) = n_i(4) = \left\lfloor \frac{2n}{5} \right\rfloor, n_i(5) = \left\lceil \frac{n-1}{5} \right\rceil$.
- If $f(u_i) = 2, n_i(4) = 1, n_i(1) = n_i(3) = \left\lfloor \frac{2n}{5} \right\rfloor, n_i(5) = \left\lceil \frac{n-1}{5} \right\rceil$.

(c) $n \bmod 5 \equiv 2$

- If $f(u_i) = 1, n_i(3) = 1, n_i(2) = n_i(4) = \frac{2n-4}{5}, n_i(5) = \frac{n+3}{5}$.
- If $f(u_i) = 2, n_i(4) = 1, n_i(1) = n_i(3) = \frac{2n-4}{5}, n_i(5) = \frac{n+3}{5}$.

(d) $n \bmod 5 \equiv 3$

- If $f(u_i) = 1, n_i(3) = 1, n_i(2) = \frac{2n-6}{5}, n_i(4) = \frac{2n-1}{5}, n_i(5) = \frac{n+2}{5}$.

- If $f(u_i) = 2, n_i(4) = 1, n_i(1) = n_i(3) = \frac{2n-1}{5}, n_i(5) = \frac{n-3}{5}$.

In all the above cases the difference between the cardinalities of the color classes does not exceed one, so our coloring is equitable.

4. $m \bmod 5 = 3$

We color first $m - 8$ copies of W_n as we have colored the corresponding vertices in Case (1). For each vertex u_i ($m - 7 \leq i \leq m$) we extend the coloring due to following conditions, dependently on n .

(a) $n \bmod 5 \equiv 0$ or $n \bmod 5 \equiv 3$

- For vertex u_{m-7} ($f(u_{m-7}) = 1$) we have $n_{m-7}(2) = 1, n_{m-7}(3) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-7}(4) = \left\lfloor \frac{2n}{5} \right\rfloor - 1, n_{m-7}(5) = \left\lceil \frac{n}{5} \right\rceil$.
- For vertex u_{m-6} ($f(u_{m-6}) = 2$) we have $n_{m-6}(1) = 1, n_{m-6}(3) = \left\lfloor \frac{2n}{5} \right\rfloor - 1, n_{m-6}(4) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-6}(5) = \left\lceil \frac{n}{5} \right\rceil$.
- For vertex u_{m-5} ($f(u_{m-5}) = 3$) we have $n_{m-5}(4) = 1, n_{m-5}(1) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-5}(2) = \left\lfloor \frac{2n}{5} \right\rfloor - 1, n_{m-5}(5) = \left\lceil \frac{n}{5} \right\rceil$.
- For vertex u_{m-4} ($f(u_{m-4}) = 4$) we have $n_{m-4}(3) = 1, n_{m-4}(1) = \left\lfloor \frac{2n}{5} \right\rfloor - 1, n_{m-4}(2) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-4}(5) = \left\lceil \frac{n}{5} \right\rceil$.
- For vertex u_{m-3} ($f(u_{m-3}) = 5$) we have $n_{m-3}(1) = 1, n_{m-3}(2) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-3}(3) = \left\lfloor \frac{2n}{5} \right\rfloor - 1, n_{m-3}(4) = \left\lceil \frac{n}{5} \right\rceil$.
- For vertex u_{m-2} ($f(u_{m-2}) = 1$) we have $n_{m-2}(2) = 1, n_{m-2}(3) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-2}(5) = \left\lfloor \frac{2n}{5} \right\rfloor - 1, n_{m-2}(4) = \left\lceil \frac{n}{5} \right\rceil$.
- For vertex u_{m-1} ($f(u_{m-1}) = 2$) we have $n_{m-1}(3) = 1, n_{m-1}(1) = \left\lfloor \frac{2n}{5} \right\rfloor - 1, n_{m-1}(5) = \left\lfloor \frac{2n}{5} \right\rfloor, n_{m-1}(4) = \left\lceil \frac{n}{5} \right\rceil$.
- For vertex u_m ($f(u_m) = 3$) we have $n_m(5) = 1, n_m(1) = \left\lfloor \frac{2n}{5} \right\rfloor, n_m(2) = \left\lfloor \frac{2n}{5} \right\rfloor - 1, n_m(4) = \left\lceil \frac{n}{5} \right\rceil$.

Each of the colors 1, 2 and 3 are used $2 + 4\lfloor 2n/5 \rfloor$ times and colors 4 and 5 are used, each one with, $2\lfloor 2n/5 \rfloor + 4\lceil n/5 \rceil + 1$ times. For $n \bmod 5 \equiv 0$ or $n \bmod 5 \equiv 3$, the difference does not exceed one.

(b) $n \bmod 5 \equiv 1$

- For vertex u_{m-7} ($f(u_{m-7}) = 1$) we have $n_{m-7}(2) = 1, n_{m-7}(3) = n_{m-7}(4) = \frac{2n-2}{5}, n_{m-7}(5) = \frac{n-1}{5}$.
- For vertex u_{m-6} ($f(u_{m-6}) = 2$) we have $n_{m-6}(1) = 1, n_{m-6}(3) = \frac{2n-2}{5}, n_{m-6}(4) = \frac{2n-7}{5}, n_{m-6}(5) = \frac{n+4}{5}$.
- For vertex u_{m-5} ($f(u_{m-5}) = 3$) we have $n_{m-5}(4) = 1, n_{m-5}(1) = n_{m-5}(2) = \frac{2n-2}{5}, n_{m-5}(5) = \frac{n-1}{5}$.
- For vertex u_{m-4} ($f(u_{m-4}) = 4$) we have $n_{m-4}(3) = 1, n_{m-4}(1) = \frac{2n-2}{5}, n_{m-4}(2) = \frac{2n-7}{5}, n_{m-4}(5) = \frac{n+4}{5}$.
- For vertex u_{m-3} ($f(u_{m-3}) = 5$) we have $n_{m-3}(1) = 1, n_{m-3}(2) = n_{m-3}(3) = \frac{2n-2}{5}, n_{m-3}(4) = \frac{n-1}{5}$.
- For vertex u_{m-2} ($f(u_{m-2}) = 1$) we have $n_{m-2}(2) = 1, n_{m-2}(3) = \frac{2n-2}{5}, n_{m-2}(5) = \frac{2n-7}{5}, n_{m-2}(4) = \frac{n+4}{5}$.
- For vertex u_{m-1} ($f(u_{m-1}) = 2$) we have $n_{m-1}(3) = 1, n_{m-1}(1) = n_{m-1}(5) = \frac{2n-2}{5}, n_{m-1}(4) = \frac{n-1}{5}$.
- For vertex u_m ($f(u_m) = 3$) we have $n_m(5) = 1, n_m(1) = \frac{2n-7}{5}, n_m(2) = \frac{2n-2}{5}, n_m(4) = \frac{n+4}{5}$.

Each of the colors 1, 2, 4 and 5 are used $(8n + 7)/5$ times and color 3 is used $(8n + 7)/5 + 1$ times.

(c) $n \bmod 5 \equiv 2$

- For vertex u_{m-7} ($f(u_{m-7}) = 1$) we have $n_{m-7}(2) = 1, n_{m-7}(3) = \frac{2n+1}{5}, n_{m-7}(4) = \frac{2n-4}{5}, n_{m-7}(5) = \frac{n-2}{5}$.
- For vertex u_{m-6} ($f(u_{m-6}) = 2$) we have $n_{m-6}(1) = 1, n_{m-6}(3) = n_{m-6}(4) = \frac{2n-4}{5}, n_{m-6}(5) = \frac{n+3}{5}$.
- For vertex u_{m-5} ($f(u_{m-5}) = 3$) we have $n_{m-5}(4) = 1, n_{m-5}(1) = \frac{2n+1}{5}, n_{m-5}(2) = \frac{2n-4}{5}, n_{m-5}(5) = \frac{n-2}{5}$.

- For vertex u_{m-4} ($f(u_{m-4}) = 4$) we have $n_{m-4}(3) = 1, n_{m-4}(1) = n_{m-4}(2) = \frac{2n-4}{5}, n_{m-4}(5) = \frac{n+3}{5}$.
- For vertex u_{m-3} ($f(u_{m-3}) = 5$) we have $n_{m-3}(1) = 1, n_{m-3}(2) = \frac{2n+1}{5}, n_{m-3}(3) = \frac{2n-4}{5}, n_{m-3}(4) = \frac{n-2}{5}$.
- For vertex u_{m-2} ($f(u_{m-2}) = 1$) we have $n_{m-2}(2) = 1, n_{m-2}(3) = n_{m-2}(5) = \frac{2n-4}{5}, n_{m-2}(4) = \frac{n+3}{5}$.
- For vertex u_{m-1} ($f(u_{m-1}) = 2$) we have $n_{m-1}(3) = 1, n_{m-1}(1) = \frac{2n-4}{5}, n_{m-1}(5) = \frac{2n+1}{5}, n_{m-1}(4) = \frac{n-2}{5}$.
- For vertex u_m ($f(u_m) = 3$) we have $n_m(5) = 1, n_m(1) = n_m(2) = \frac{2n-4}{5}, n_m(4) = \frac{n+3}{5}$.

Each of the colors 1, 2, 3 and 5 are used $(8n+4)/5 + 1$ times and color 4 is used $(8n+4)/5$ times.

(d) $n \bmod 5 \equiv 4$

- For vertex u_{m-7} ($f(u_{m-7}) = 1$) we have $n_{m-7}(2) = 1, n_{m-7}(3) = n_{m-7}(4) = \frac{2n-3}{5}, n_{m-7}(5) = \frac{n+1}{5}$.
- For vertex u_{m-6} ($f(u_{m-6}) = 2$) we have $n_{m-6}(1) = 1, n_{m-6}(3) = n_{m-6}(4) = \frac{2n-3}{5}, n_{m-6}(5) = \frac{n+1}{5}$.
- For vertex u_{m-5} ($f(u_{m-5}) = 3$) we have $n_{m-5}(4) = 1, n_{m-5}(1) = n_{m-5}(2) = \frac{2n-3}{5}, n_{m-5}(5) = \frac{n+1}{5}$.
- For vertex u_{m-4} ($f(u_{m-4}) = 4$) we have $n_{m-4}(3) = 1, n_{m-4}(1) = n_{m-4}(2) = \frac{2n-3}{5}, n_{m-4}(5) = \frac{n+1}{5}$.
- For vertex u_{m-3} ($f(u_{m-3}) = 5$) we have $n_{m-3}(1) = 1, n_{m-3}(2) = n_{m-3}(3) = \frac{2n-3}{5}, n_{m-3}(4) = \frac{n+1}{5}$.
- For vertex u_{m-2} ($f(u_{m-2}) = 1$) we have $n_{m-2}(2) = 1, n_{m-2}(3) = n_{m-2}(5) = \frac{2n-3}{5}, n_{m-2}(4) = \frac{n+1}{5}$.
- For vertex u_{m-1} ($f(u_{m-1}) = 2$) we have $n_{m-1}(3) = 1, n_{m-1}(1) = n_{m-1}(5) = \frac{2n-3}{5}, n_{m-1}(4) = \frac{n+1}{5}$.
- For vertex u_m ($f(u_m) = 3$) we have $n_m(5) = 1, n_m(1) = n_m(2) = \frac{2n-3}{5}, n_m(4) = \frac{n+1}{5}$.

Each of the colors are used $(8n + 8)/5$ times.

In all the above cases the difference between the cardinalities of the color classes does not exceed one, so our coloring is equitable.

5. $m \bmod 5 \equiv 4$

We color first $m - 4$ copies of W_n as we have colored the corresponding vertices in Case (1). Then, we color last four copies in the following way. For each vertex u_i , ($m - 3 \leq i \leq m$), we color the corresponding copy of W_n due the following conditions, dependently on n .

(a) $n \bmod 5 \equiv 0$

If $f(u_i) = j$, $1 \leq j \leq 4$, then

- $n_i((j + 1) \bmod 4) = 1$,
- $n_i((j + 2) \bmod 4) = \frac{2n}{5}$,
- $n_i((j + 3) \bmod 4) = \frac{2n - 5}{5}$,
- $n_i(5) = \frac{n}{5}$.

(b) $n \bmod 5 \equiv 1$

For vertex u_{m-3} ($f(u_{m-3}) = 1$) we have $n_{m-3}(2) = 1, n_{m-3}(3) = \frac{2n - 2}{5}$,

$$n_{m-3}(4) = \frac{2n - 7}{5}, n_{m-3}(5) = \frac{n + 4}{5}.$$

For vertices u_i , $m - 2 \leq i \leq m$, if $f(u_i) = j$, $1 \leq j \leq 4$, then

- $n_i((j + 1) \bmod 4) = 1$,
- $n_i((j + 2) \bmod 4) = n_i((j + 3) \bmod 4) = \frac{2n - 2}{5}$,
- $n_i(5) = \frac{n - 1}{5}$.

(c) $n \bmod 5 \equiv 2$

For vertices u_i , $m - 3 \leq i \leq m - 2$, if $f(u_i) = j$, $1 \leq j \leq 2$, then

- $n_i((j + 1) \bmod 4) = 1$,
- $n_i((j + 2) \bmod 4) = n_i((j + 3) \bmod 4) = \frac{2n - 4}{5}$,
- $n_i(5) = \frac{n + 3}{5}$.

For vertices u_i , $m - 1 \leq i \leq m$, if $f(u_i) = j$, $3 \leq j \leq 4$, then

- $n_i((j + 1) \bmod 4) = 1$,
- $n_i((j + 2) \bmod 4) = \frac{2n + 1}{5}$,
- $n_i((j + 3) \bmod 4) = \frac{2n - 4}{5}$,
- $n_i(5) = \frac{n - 2}{5}$.

(d) $n \bmod 5 \equiv 3$

If $f(u_i) = j$, $1 \leq j \leq 4$, then

- $n_i((j + 1) \bmod 4) = 1,$
- $n_i((j + 2) \bmod 4) = \frac{2n - 1}{5},$
- $n_i((j + 3) \bmod 4) = \frac{2n - 6}{5},$
- $n_i(5) = \frac{n + 2}{5}.$

(e) $n \bmod 5 \equiv 4$

If $f(u_i) = j, 1 \leq j \leq 4,$ then

- $n_i((j + 1) \bmod 4) = 1,$
- $n_i((j + 2) \bmod 4) = n_i((j + 3) \bmod 4) = \frac{2n - 3}{5},$
- $n_i(5) = \frac{n + 1}{5}.$

In all the above cases the difference between the cardinalities of the color classes does not exceed one, so our coloring is equitable. Hence $\chi_{=}(G \circ W_n) \leq 5.$ By the definition of corona graph for each vertex u_i of $G,$ there exists a copy of W_n whose vertices are adjacent to the vertex $u_i.$

Case 1: If $m \bmod 2 \equiv 1, n \geq 7$

In this case either both m and n are odd (or) m is odd and n is even.

(a) If m and n are odd.

Since $\chi(W_n) = 3$ for odd $n,$ we need at least 4 colors for coloring each copy of W_n and the corresponding vertex of $G.$ In this coloring, since m is odd there exists atleast one color which reappears in $\langle \{u_i : 1 \leq i \leq m\} \rangle.$ Let the color $j (1 \leq j \leq 4)$ reappears at the vertex $u_i (5 \leq i \leq m).$ Then the center vertex of the copy W_n corresponding to the vertex $u_i,$ receives a color $k (1 \leq k \leq 4),$ where $k \neq j.$ Other vertices of W_n receive the colors other than j and $k.$ (i.e) The number of possible colors to color these vertices is two. Hence it is clear that for the case of $n \geq 5,$ it is not possible to color the vertices of the cycle C_{n-1} of W_n equitably with two colors. Therefore $\chi_{=}(G \circ W_n) \geq 5.$ Hence $\chi_{=}(G \circ W_n) = 5$ for m and n are odd.

(b) If m is odd and n is even.

Since $\chi(W_n) = 4$ for even $n,$ the graph $G \circ W_n$ requires at least 5 colors. Hence $\chi_{=}(G \circ W_n) = 5$ for m is odd and n is even.

Case 2: If m and n are even, $n \geq 4$

Since $\chi(W_n) = 4$ for even $n,$ graph $G \circ W_n$ requires at least 5 colors. Therefore $\chi_{=}(G \circ W_n) \geq 5.$

Hence $\chi_{=}(G \circ W_n) = 5$ for even $n.$

□

Next, we consider coronas, where the set of vertices of graph G includes exactly three elements.

Theorem 2.5. *Let G be an equitably 3-colorable graph with $m = 3$ vertices. Then*

1. $\chi_=(G \circ W_5) = 4$.
2. $\chi_=(G \circ W_n) = 5$ $n = 7, 9, 11, 13, 15, 17$.
3. $\chi_=(G \circ W_n) = 5$ $n \geq 19$, if n is odd.
4. $\chi_=(G \circ W_n) = 5$ $n = 4, 6, 8, 10$.
5. $\chi_=(G \circ W_n) = 6$ $n \geq 12$, if n is even.

Proof. Let $\{u_i : 1 \leq i \leq 3\}$ be the set of vertices of G .

1. We color $G \circ W_5$ as for the following procedure.

- For vertex u_1 ($f(u_1) = 1$) we have $n_1(2) = 1, n_1(3) = n_1(4) = 2$.
- For vertex u_2 ($f(u_2) = 2$) we have $n_2(3) = 1, n_2(4) = n_2(1) = 2$.
- For vertex u_3 ($f(u_3) = 3$) we have $n_3(4) = 1, n_3(1) = n_3(2) = 2$.

In the above cases the difference between the cardinalities of the color classes does not exceed one, so our coloring is equitable. Hence $\chi_=(G \circ W_5) \leq 4$. Since W_5 is 3-colorable, at each copy of W_5 of $G \circ W_5$, there exists one more color. Therefore $\chi_=(G \circ W_5) \geq 4$. hence $\chi_=(G \circ W_5) = 4$.

2. Assign the color i to the vertex u_i ($1 \leq i \leq 3$), color 4 to the vertex u_{1n} , color 5 to the vertex u_{2n} and color 1 to the vertex u_{3n} . Since C_{n-1} is of even order, we require only two colors for proper coloring of C_{n-1} . We use three colors in each C_{n-1} of W_n in $G \circ W_n$. We use the colors 2,3,5 to the vertices of C_{n-1} of W_n at u_1 . Similarly we use the colors 1,3,4 and 4,5,2 to the vertices of C_{n-1} of W_n at u_2 and u_3 respectively. The number of appearance of the colors are given in the following cases.

(a) $n = 7, 17$

- For vertex u_1 ($f(u_1) = 1$) we have $n_1(4) = 1, n_1(2) = \frac{2n+1}{5}, n_1(3) = \frac{2n-4}{5}, n_1(5) = \frac{n-2}{5}$.
- For vertex u_2 ($f(u_2) = 2$) we have $n_2(5) = 1, n_2(1) = \frac{n-1}{2}, n_2(3) = \left\lceil \frac{n-1}{4} \right\rceil, n_2(4) = \left\lfloor \frac{n-1}{4} \right\rfloor$.
- For vertex u_3 ($f(u_3) = 3$) we have $n_3(1) = 1, n_3(4) = \frac{2n-4}{5}, n_3(5) = \frac{2n+1}{5}, n_3(2) = \frac{n-2}{5}$.

(b) $n = 9$

- For vertex u_1 ($f(u_1) = 1$) we have $n_1(4) = 1, n_1(2) = n_1(3) = 3, n_1(5) = 2$.
- For vertex u_2 ($f(u_2) = 2$) we have $n_2(5) = 1, n_2(1) = 4, n_2(3) = n_2(4) = 2$.

- For vertex u_3 ($f(u_3) = 3$) we have $n_3(1) = 1, n_3(4) = n_3(5) = 3, n_3(2) = 2$.
- (c) $n = 11$
- For vertex u_1 ($f(u_1) = 1$) we have $n_1(4) = 1, n_1(2) = n_1(3) = 4, n_1(5) = 2$.
 - For vertex u_2 ($f(u_2) = 2$) we have $n_2(5) = 1, n_2(1) = 5, n_2(3) = 3, n_2(4) = 2$.
 - For vertex u_3 ($f(u_3) = 3$) we have $n_3(1) = 1, n_3(4) = n_3(5) = 4, n_3(2) = 2$.
- (d) $n = 13$
- For vertex u_1 ($f(u_1) = 1$) we have $n_1(4) = 1, n_1(2) = n_1(3) = 5, n_1(5) = 2$.
 - For vertex u_2 ($f(u_2) = 2$) we have $n_2(5) = 1, n_2(1) = 6, n_2(3) = 3, n_2(4) = 3$.
 - For vertex u_3 ($f(u_3) = 3$) we have $n_3(1) = 1, n_3(4) = n_3(5) = 5, n_3(2) = 2$.
- (e) $n = 15$
- For vertex u_1 ($f(u_1) = 1$) we have $n_1(4) = 1, n_1(2) = n_1(3) = 6, n_1(5) = 2$.
 - For vertex u_2 ($f(u_2) = 2$) we have $n_2(5) = 1, n_2(1) = 7, n_2(3) = 3, n_2(4) = 4$.
 - For vertex u_3 ($f(u_3) = 3$) we have $n_3(1) = 1, n_3(4) = 4, n_3(5) = 7, n_3(2) = 2$.

In the above cases the difference between the cardinalities of the color classes does not exceed one, so our coloring is equitable. Hence $\chi_=(G \circ W_n) \leq 5$.

Since G is 3-colorable, let i be the color assigned to the vertex u_i ($1 \leq i \leq 3$) of $G \circ W_n$. Let j ($1 \leq j \leq 4$), ($i \neq j$) be the color assigned to the center vertices of each copy W_n of $G \circ W_n$. The other vertices of these copies receive the colors other than i and j . (i.e) The number of possible colors to color these vertices is two. Hence it is clear that for the case of $n = 7, 9, 11, 13, 15, 17$, it is not possible to color the vertices of the cycle C_{n-1} of W_n equitably with two colors. Therefore $\chi_=(G \circ W_n) \geq 5$. Hence $\chi_=(G \circ W_n) = 5$ for $n = 7, 9, 11, 13, 15, 17$.

3. Suppose that $G \circ W_n$ is 4-equitably colorable. Since G is 3-colorable, let it be colored by the color 1,2 and 3. Let u_i receives the color i ($1 \leq i \leq 3$). Then u_{1n}, u_{2n} and u_{3n} should receive any two of the three color 1,2,3 and the color 4.

Let u_{1n} receive 4, u_{2n} receive 1 and u_{3n} receive 2. Then u_{1i} ($1 \leq i \leq n-1$) receives the color 2, $\frac{n-1}{2}$ times and 3, $\frac{n-1}{2}$ times. u_{2i} ($1 \leq i \leq n-1$) receives the color 3, $\frac{n-1}{2}$ times, the color 4, $\frac{n-1}{2}$ times. Similarly u_{3i} receives the color 1, $\frac{n-1}{2}$ times and the color 1, $\frac{n-1}{2}$ times.

Number of appearance of each colors 1 and 2 are, $\frac{n+3}{2}$ times respectively and number of appearance of each colors 3 and 4 are, n times respectively.

As the above mentioned partition does not imply the equitable partition, it is concluded that $G \circ W_n$ should not be equitable 4-colorable.

Hence $\chi_=(G \circ W_n) \geq 5$

Suppose that $G \circ W_n$ is 5-equitable colorable. Let G be colored by the colors 1, 2 and 3. Let u_i receives the color i ($1 \leq i \leq 3$). Since $G \circ W_n$ is 5-equitable colorable, any two of the vertices u_{1n}, u_{2n} and u_{3n} receives the color 4 and 5 (Say u_{1n}, u_{2n}) and remaining vertex u_{3n} should receive the color 1.

For the case of $n \geq 19$, if we use the above coloring with 5 colors, then the maximum of appearance of color 1, $\frac{n-1}{2} + 2 = \frac{n+3}{2}$ times.

Remaining number of vertices to be colored are, $3n + 3 - \frac{n+3}{2} = \frac{5n+3}{2}$.

Number of vertices which receive each colors of 2, 3, 4 and 5 are $\frac{\frac{5n+3}{2}}{4} = \frac{5n+3}{8}$.

For $n \geq 19$, $\left\lceil \frac{5n+3}{2} \right\rceil - \left\lceil \frac{n+3}{2} \right\rceil \geq 2$.

(i.e) it may not be possible to equitably color $G \circ W_n$ with 5 colors.

$\chi_=(G \circ W_n) \geq 6$.

- For vertex $u_1 (f(u_1) = 1)$ we have $n_1(4) = 1, n_1(2) = n_1(3) = \frac{n-1}{2}$.
- For vertex $u_2 (f(u_2) = 2)$ we have $n_2(5) = 1, n_2(1) = n_2(6) = \frac{n-1}{2}$.
- For vertex $u_3 (f(u_3) = 3)$ we have $n_3(6) = 1, n_3(5) = n_3(4) = \frac{n-1}{2}$.

In the above cases the difference between the cardinalities of color classes does not exceed one, so our coloring is equitable. Hence $\chi_=(G \circ W_n) = 6, n \geq 19$, if n is odd.

4. Since n is even W_n has odd cycle C_{n-1} . Minimum number of colors assigned to color any cycle is 3. Hence $u_{in} (1 \leq i \leq n)$ should have a fourth color and hence $u_i (1 \leq i \leq n)$ must receive a fifth color. Hence $\chi_=(G \circ W_n) \geq 5$.

Now we partition the vertex set $V(G \circ W_n)$ as follows,

$$\begin{aligned} V_1 &= \{u_1, u_{21}, u_{23}, u_{25}, u_{28}, u_{3n}\} \\ V_2 &= \{u_2, u_{11}, u_{14}, u_{18}, u_{33}, u_{36}, u_{39}\} \\ V_3 &= \{u_3, u_{12}, u_{15}, u_{17}, u_{24}, u_{27}\} \\ V_4 &= \{u_{1n}, u_{22}, u_{26}, u_{29}, u_{32}, u_{35}, u_{37}\} \\ V_5 &= \{u_{2n}, u_{13}, u_{16}, u_{19}, u_{31}, u_{34}, u_{38}\} \end{aligned}$$

Clearly V_1, V_2, V_3, V_4 and V_5 are independent set of $G \circ W_n$. Hence $||V_i| - |V_j|| \leq 1$ for every $i \neq j$. Hence $\chi_=(G \circ W_n) = 5, 4 \leq n \leq 10$, if n is even.

5. Let $n_i(k)$ be the number of appearance of the color k in the copy of W_n corresponding to the vertex u_i of G in $G \circ W_n$.

Let $f(u_i) = j$ be the color assigned to each vertices $u_i (1 \leq i \leq m)$ of G . Since G is 6-colorable j takes the values in the range $1 \leq j \leq 6$.

- For vertex $u_1 (f(u_1) = 1)$ we have $n_1(2) = n_1(5) = 1, n_1(3) = n_1(4) = \frac{n-2}{2}$.
- For vertex $u_2 (f(u_2) = 2)$ we have $n_2(3) = n_2(1) = 1, n_2(5) = n_2(6) = \frac{n-2}{2}$.
- For vertex $u_3 (f(u_3) = 3)$ we have $n_3(6) = n_3(4) = 1, n_3(1) = n_3(2) = \frac{n-2}{2}$.

In the above cases the difference between the cardinalities of the color classes does not exceed one, so our coloring is equitable. Hence $\chi_=(G \circ W_n) \leq 6$

Since n is even, we require at least 3 colors to color each C_{n-1} of W_n , one color for the centre vertex of W_n and one color corresponding to the vertex of G . Hence we may assume that $\chi_=(G \circ W_n) = 5$. It is clear that one of these five colors appears twice in $\langle \{u_i : 1 \leq i \leq 3\} \cup \{u_{in} : 1 \leq i \leq 3\} \rangle$, let it be color j ($1 \leq j \leq 5$). This color j can be assigned only $\frac{n-2}{2}$ times in any of the C_{n-1} copy of W_{n-1} . This violate the equitable conclusion.

Therefore $\chi_=(G \circ W_n) \geq 6$. Hence $\chi_=(G \circ W_n) = 6$. □

3. Conclusion

We notice that the results can be extended into further products of graphs.

Corollary 3.1. *Let G be an equitably 4-colorable graph on, $m \geq 2$, vertices, let m is even, n is odd, and $n \geq 4$, and $l \geq 1$. Then*

$$\chi_=(G \circ W_n) = 4.$$

Proof. We use the principle of mathematical induction due to number l .

1. $l=1$

The truth follows immediately from Theorem 2.2.

2. Induction hypothesis for l . It means that $\chi_=(G \circ^l W_n) = 4$ for n odd and $m = |V(G)|$ even.

3. We must show that $\chi_=(G \circ^{l+1} W_n) = 4$ for graphs under consideration.

Let us notice that graph from induction hypothesis $G \circ^l W_n$ is an equitably 4-colorable graph, it means a graph fulfilling the assumption of Theorem 2.2. Its number of vertices, equals to $m(n+1)^l$ is an even number for m even. So, $\chi_=(G \circ^{l+1} W_n) = 4$. □

Corollary 3.2. *Let G be an equitably 5-colorable graph on m vertices and let $m \geq 2$, $n \geq 4$, $l \geq 1$. Then*

$$\chi_=(G \circ^l W_n) = \begin{cases} = 5 & \text{for } n \text{ even,} \\ \leq 5 & \text{for } m \text{ and } n \text{ odd.} \end{cases}$$

Proof. Follows immediately from Theorem 2.4. □

Acknowledgement

With due Respect, the authors sincerely thank the referee for his careful reading, excellent comments and fruitful suggestions that have resulted in the improvement of the quality of this manuscript.

References

- [1] B. L. Chen and K. W. Lih, Equitable coloring of trees, *J. Combin. Theory Ser. B* **61** (1994), 83–87.
- [2] R. Frucht and F. Harary, On the corona of two graphs, *Aequationes Math.* **4** (1970), 322–325.
- [3] H. Furmańczyk, Equitable coloring of graph products, *Opuscula Mathematica* **26** (2006), 31–44.
- [4] H. Furmańczyk, K. Kaliraj, M. Kubale, and J. Vernold Vivin, Equitable coloring of corona product of graphs, *Advances and Applications of Discrete Mathematics* **11**(2) (2013), 103–120.
- [5] S. Klavzar, R. Hammack, and W. Imrich, *Handbook of graph products*, CRC Press (2011).
- [6] M. Kubale, *Graph Colorings*, American Mathematical Society Providence, Rhode Island- (2004).
- [7] K. W. Lih and P. L. Wu, On equitable coloring of bipartite graphs, *Discrete Math.* **151** (1996), 155–160.
- [8] W. H. Lin and G. J. Chang, Equitable colorings of Cartesian products of graphs, *Disc. App. Math.* **160** (2012), 239–247.
- [9] W. Meyer, Equitable coloring, *Amer. Math. Monthly* **80** (1973), 920–922.
- [10] W. Wang and K. Zhang, Equitable colorings of line graphs and complete r -partite graphs, *Systems Science and Mathematical Sciences* **13** (2000), 190–194.
- [11] H. P. Yap and Y. Zhang, The Equitable Δ -Coloring Conjecture holds for outerplanar graphs, *Bulletin of the Inst. of Math. Academia Sinica* **25** (1997), 143–149.
- [12] H. P. Yap and Y. Zhang, Equitable colourings of planar graphs, *J. Comb. Math. Comb. Comput.* **27** (1998), 97–105.