



Note on Parity Factors of Regular Graphs

Hongliang Lu^a, Yuqing Lin^b

^aDepartment of Mathematics

Xi'an Jiaotong University, Xi'an 710049, PR China

^bSchool of Electrical Engineering and Computer Science,
The University of Newcastle, Australia

luhongliang215@sina.com, yuqing.lin@newcastle.edu.au

Abstract

In this paper, we obtain a sufficient condition for the existence of parity factors in a regular graph in terms of edge-connectivity. Moreover, we also show that our condition is sharp.

Keywords: Parity Factors, edge-connectivity

Mathematics Subject Classification : 05C70

1. Preliminaries

Let $G = (V, E)$ be a graph with vertex set $V(G)$ and edge set $E(G)$. The number of vertices of a graph G is called the *order* of G and is denoted by n . The number of edges of G is called the *size* of G and is denoted by e . For a vertex v of graph G , the number of edges of G incident to v is called the *degree* of v in G and is denoted by $d_G(v)$. For two subsets $S, T \subseteq V(G)$, let $e_G(S, T)$ denote the number of edges of G joining S to T .

Let H be a function associating a subset of \mathbb{Z} to each vertex of G . A spanning subgraph F of graph G is called an *H -factor* of G if

$$d_F(x) \in H(x) \quad \text{for every vertex } x \in V(G). \quad (1)$$

For a spanning subgraph F of G and for a vertex v of G , define

$$\delta(H; F, v) = \min\{|d_F(v) - i| \mid i \in H_v\},$$

Received: 6 February 2013, Revised: 8 April 2013, Accepted: 24 April 2013.

and let $\delta(H; F) = \sum_{x \in V(G)} \delta(H; F, x)$. Thus a spanning subgraph F is an H -factor if and only if $\delta(H; F) = 0$. Let

$$\delta_H(G) = \min\{\delta(H; F) \mid F \text{ are spanning subgraphs of } G\}.$$

A spanning subgraph F is called H -optimal if $\delta(H; F) = \delta_H(G)$. The H -factor problem is to determine the value $\delta_H(G)$. An integer h is called a *gap* of $H(v)$ if $h \notin H(v)$ but $H(v)$ contains an element less than h and an element greater than h . Lovász [11] gave a structural description on the H -factor problem in the case where $H(v)$ has no two consecutive gaps for all $v \in V(G)$ and showed that the problem is NP-complete without this restriction. Moreover, he also conjectured that the decision problem of determining whether a graph has an H -factor is polynomial in the case where $H(v)$ has no two consecutive gaps for all $v \in V(G)$. Cornuéjols [5] proved the conjecture.

Let therefore $g, f : V \rightarrow Z^+$ such that $g(v) \leq f(v)$ and $g(v) \equiv f(v) \pmod{2}$ for every $v \in V$. Then a spanning subgraph F of G is called a (g, f) -parity-factor, if $g(v) \leq d_F(v) \leq f(v)$ and $d_F(v) \equiv f(v) \pmod{2}$ for all $v \in V$. Clearly, a (g, f) -parity-factor is a special kind of H -factor and it has been shown that the decision problem of determining whether a graph has a (g, f) -parity factor is polynomial.

Let a, b be two integers such that $1 \leq a \leq b$ and $a \equiv b \pmod{2}$. If $g(v) = a$ and $f(v) = b$ for all $v \in V(G)$, then a (g, f) -parity-factor is called an (a, b) -parity factor. Let $n \geq 1$ be odd. If $a = 1$ and $b = n$, then an (a, b) -parity factor is called a $(1, n)$ -odd factor. There is also a special case of the (g, f) -factor problem which is called the *even factor problem*, i.e., the problem with $g(v) = 2$, $f(v) \geq |V(G)|$ and $f(v) \equiv g(v) \pmod{2}$ for all $v \in V(G)$.

Fleischner gave a sufficient condition for a graph to have an even factor in terms of edge connectivity.

Theorem 1.1 (Fleischner,[8]; Lovász, [12]). *If G is a bridgeless graph with $\delta(G) \geq 3$, then G has an even factor.*

For a general graph G and an integer k , a spanning subgraph F such that

$$d_F(x) = k \text{ for all } x \in V(G)$$

is called a k -factor. In fact, a k -factor is also a (k, k) -parity factor.

The first investigation of the $(1, n)$ -odd factor problem is due to Amahashi [2], who gave a Tutte type characterization for graphs having a global odd factor.

Theorem 1.2 (Amahashi). *Let n be an odd integer. A graph G has a $(1, n)$ -odd factor if and only if*

$$o(G - S) \leq n |S| \quad \text{for all subsets } S \subset V(G). \quad (2)$$

For general odd value functions h , Cui and Kano [6] established a Tutte type of theorem.

Theorem 1.3 (Cui and Kano, [6]). *Let $h : V(G) \rightarrow N$ be odd value function. A graph G has a $(1, h)$ -odd factor if and only if*

$$o(G - S) \leq h(S) \quad \text{for all subsets } S \subset V(G). \quad (3)$$

Now there are many results on consecutive factors (i.e. (g, f) -factor). But the research progress on non-consecutive factors is slow. In non-consecutive factor problems, (g, f) -parity factors have many similar properties with k -factors. So we believe that many results on k -factors can be extended to (g, f) -factor. In this paper, we will extend a result on k -factors of regular graphs to the (g, f) -parity-factors.

Now let us recall one of the classical results due to Petersen.

Theorem 1.4 (Petersen [13]). *Let r and k be integers such that $1 \leq k \leq r$. Every $2r$ -regular graph has a $2k$ -factor.*

Considering the edge-connectivity, Gallai [7] proved the following result.

Theorem 1.5 (Gallai [7]). *Let r and k be integers such that $1 \leq k < r$, and G an m -edge-connected r -regular graph, where $m \geq 1$. If one of the following conditions holds, then G has a k -factor.*

- (i) r is even, k is odd, $|G|$ is even, and $\frac{r}{m} \leq k \leq r(1 - \frac{1}{m})$;
- (ii) r is odd, k is even and $2 \leq k \leq r(1 - \frac{1}{m})$;
- (iii) r and k are both odd and $\frac{r}{m} \leq k$.

Bollobás, Satio and Wormald [3] improved above the result.

Theorem 1.6 (Bollobás, Saito and Wormald). *Let r and k be integers such that $1 \leq k < r$, and G be an m -edge-connected r -regular graph, where $m \geq 1$ is a positive integer. Let $m^* \in \{m, m+1\}$ such that $m^* \equiv 1 \pmod{2}$. If one of the the following conditions holds, then G has a k -factor.*

- (i) r is odd, k is even and $2 \leq k \leq r(1 - \frac{1}{m^*})$;
- (ii) r and k are both odd and $\frac{r}{m^*} \leq k$.

In this paper, we extend Theorems 1.5 and 1.6 to (a, b) -factors. The main tool in our proofs is the following theorem of Lovász (see[11]).

Theorem 1.7 (Lovász [11]). *G has a (g, f) -parity factor if and only if for all disjoint subsets S and T of $V(G)$,*

$$\delta(S, T) = f(S) + \sum_{x \in T} d_G(x) - g(T) - e_G(S, T) - \tau \geq 0,$$

where τ denotes the number of components C , called f -odd components of $G - (S \cup T)$ such that $e_G(V(C), T) + f(V(C)) \equiv 1 \pmod{2}$. Moreover, $\delta(S, T) \equiv f(V(G)) \pmod{2}$.

2. Main Theorem

Theorem 2.1. *Let a, b and r be integers such that $1 \leq a \leq b < r$ and $a \equiv b \pmod{2}$. Let G be an m -edge-connected r -regular graph with n vertices. Let $m^* \in \{m, m + 1\}$ such that $m^* \equiv 1 \pmod{2}$. If one of the following conditions holds, then G has an (a, b) -parity factor.*

- (i) r is even, a, b are odd, $|G|$ is even, $\frac{r}{m} \leq b$ and $a \leq r(1 - \frac{1}{m})$;
- (ii) r is odd, a, b are even and $a \leq r(1 - \frac{1}{m^*})$;
- (iii) r, a, b are odd and $\frac{r}{m^*} \leq b$.

By Theorem 1.6, (ii) and (iii) are true. Now we prove (i). Let $\theta_1 = \frac{a}{r}$ and $\theta_2 = \frac{b}{r}$. Then $0 < \theta_1 \leq \theta_2 < 1$. Suppose that G contains no (a, b) -parity factors. By Theorem 1.7, there exist two disjoint subsets S and T of $V(G)$ such that $S \cup T \neq \emptyset$, and

$$-2 \geq \delta(S, T) = b|S| + \sum_{x \in T} d_G(x) - a|T| - e_G(S, T) - \tau, \tag{4}$$

where τ is the number of a -odd (i.e. b -odd) components C of $G - (S \cup T)$. Let C_1, \dots, C_τ denote a -odd components of $G - S - T$ and $D = C_1 \cup \dots \cup C_\tau$.

Note that

$$\begin{aligned} -2 \geq \delta(S, T) &= b|S| + \sum_{x \in T} d_G(x) - a|T| - e_G(S, T) - \tau \\ &= b|S| + (r - a)|T| - e_G(S, T) - \tau \\ &= \theta_2 r|S| + (1 - \theta_1)r|T| - e_G(S, T) - \tau \\ &= \theta_2 \sum_{x \in S} d_G(x) + (1 - \theta_1) \sum_{x \in T} d_G(x) - e_G(S, T) - \tau \\ &\geq \theta_2 (e_G(S, T) + \sum_{i=1}^{\tau} e_G(S, C_i)) + (1 - \theta_1) (e_G(S, T) + \sum_{i=1}^{\tau} e_G(T, C_i)) - e_G(S, T) - \tau \\ &= \sum_{i=1}^{\tau} (\theta_2 e_G(S, C_i) + (1 - \theta_1) e_G(T, C_i) - 1) + (\theta_2 - \theta_1) e_G(S, T) \\ &\geq \sum_{i=1}^{\tau} (\theta_2 e_G(S, C_i) + (1 - \theta_1) e_G(T, C_i) - 1). \end{aligned}$$

Since G is connected and $0 < \theta_1 \leq \theta_2 < 1$, so $\theta_2 e_G(S, C_i) + (1 - \theta_1) e_G(T, C_i) > 0$ for each C_i . Hence we will obtain a contradiction by showing that for every $C = C_i, 1 \leq i \leq \tau$, we have

$$\theta_2 e_G(S, C) + (1 - \theta_1) e_G(T, C) \geq 1. \tag{5}$$

These inequalities imply

$$\begin{aligned} -2 \geq \delta_G(S, T) &\geq \sum_{i=1}^{\tau} (\theta_2 e_G(S, C_i) + (1 - \theta_1) e_G(T, C_i) - 1) \\ &> \sum_{i=1}^{\tau-2} (\theta_2 e_G(S, C_i) + (1 - \theta_1) e_G(T, C_i) - 1) - 2 \geq -2, \end{aligned}$$

which is impossible.

Now, we will prove the 5 is true. Since C is an a -odd component of $G - (S \cup T)$, we have

$$a|C| + e_G(T, C) \equiv 1 \pmod{2}. \tag{6}$$

Moreover, since

$$r|C| = \sum_{x \in V(C)} d_G(x) = e_G(S \cup T, C) + 2|E(C)|,$$

we have

$$r|C| = e_G(S \cup T, C) \pmod{2}. \tag{7}$$

It is obvious that the two inequalities $e_G(S, C) \geq 1$ and $e_G(T, C) \geq 1$ imply

$$\theta_2 e_G(S, C) + (1 - \theta_1) e_G(T, C) \geq \theta_2 + 1 - \theta_1 = 1.$$

Hence we may assume $e_G(S, C) = 0$ or $e_G(T, C) = 0$.

We consider the condition (i). If $e_G(S, C) = 0$, then $e_G(T, C) \geq m$. Since $a \leq r(1 - \frac{1}{m})$, then $\theta_1 \leq 1 - \frac{1}{m}$ and so $1 \leq (1 - \theta_1)m$. By substituting $e_G(T, C) \geq m$ and $e_G(S, C) = 0$ into (5), we have

$$(1 - \theta_1) e_G(T, C) \geq (1 - \theta_1)m \geq 1.$$

If $e_G(T, C) = 0$, then $e_G(S, C) \geq m$. Since $\frac{r}{m} \leq b$, hence $\theta_2 m \geq 1$, and so we obtain

$$\theta_2 e_G(S, C) \geq \theta_2 m \geq 1.$$

Consequently, condition (i) guarantees (5) holds and thus (i) is true. The proof is completed. \square

Remark: The edge connectivity conditions in Theorem 2.1 are sharp.

We will give the construction for condition (i) of Theorem 2.1. For (ii) and (iii), the constructions are similar. Let $r \geq 2$ be an even integer, $a, b \geq 1$ two odd integers and $2 \leq m \leq r - 2$ an even integer such that $b < r/m$ or $r(1 - \frac{1}{m}) < a$. Since G has an (a, b) -parity factor if and only if G has an $(r - b, r - a)$ -parity factor, so we can assume $b < r/m$. Let $J(r, m)$ be the complete graph K_{r+1} from which a matching of size $m/2$ is deleted. Take r disjoint copies of $J(r, m)$. Add m new vertices and connect each of these vertices to a vertex of degree $r - 1$ of $J(r, m)$. This gives an m -edge-connected r -regular graph denoted by G . Let S denote the set of m new vertices and $T = \emptyset$. Let τ denote the number of components C , which are called a -odd components of $G - (S \cup T)$ and $e_G(V(C), T) + a|C| \equiv 1 \pmod{2}$. Then we have $\tau = r$, and

$$\delta(S, T) = b|S| + \sum_{x \in T} d_{G-S}(x) - a|T| - \tau(S, T) = bm - r < 0.$$

So by Theorem 1.7, G contains no (a, b) -parity factors.

Acknowledgement

This work was supported by the National Natural Science Foundation of China (No. 11101329)

- [1] J. Akiyama and M. Kano, Factors and factorizations of graphs-a survey, *J. Graph Theory*, **9** (1985), 1-42.
- [2] A. Amahashi, On factors with all degree odd, *Graphs and Combin.*, **1** (1985), 111–114.
- [3] B. Bollobás, A. Satio, and N. C. Wormald, Regular factors of regular graphs, *J. Graph Theory*, **9** (1985), 97-103.
- [4] L. Collatz and U. Sinogowitz, Spektren endlicher Grafen, *Abh. Math. Sem. Univ. Hamburg*, **99** (2009), 287-297.
- [5] G. Cornuéjols, General factors of graphs, *J. Combin. Theory Ser. B*, **45** (1988), 185-198.
- [6] Y. Cui and M. Kano, Some results on odd factors of graphs, *J. Graph Theory*, **12** (1988), 327–333.
- [7] T. Gallai, The factorisation of graphs, *Acta Math. Acad. Sci. Hung.*, **1** (1950), 133-153.
- [8] H. Fleischner, Spanning Eulerian subgraphs, the Splitting Lemma, and Petersen’s Theorem, *Discrete Math.*, **101** (1992), 33–37.
- [9] C. Godsil and G. Royle, Algebraic Graph Theory, Springer Verlag New York, (2001).
- [10] M. Kano, $[a, b]$ -factorization of a graph. *J. Graph Theory*, **9** (1985), 129-146.
- [11] L. Lovász, The factorization of graphs, II, *Acta Math. Sci. Hungar.*, **23** (1972), 223-246.
- [12] L. Lovász, Combinatorial Problems and Exercises, North-Holland, Amsterdam (1979).
- [13] J. Petersen, Die Theorie der regulären Graphen, *Acta Math.*, **15** (1891), 193-220.
- [14] W. T. Tutte, The factors of graphs, *Canad. J. Math.*, **4** (1952), 314-328.